

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
19 February 2004 (19.02.2004)

PCT

(10) International Publication Number
WO 2004/014371 A1

- (51) International Patent Classification⁷: **A61K 31/425**, C07D 277/82, 493/04, 417/12, A61P 31/18
- (74) Agent: **DAELEMANS, Frank**; Tibotec-Virco Comm. VA, Generaal De Wittelaan L11 B3, B-2800 Mechelen (BE).
- (21) International Application Number: **PCT/EP2003/050359**
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 4 August 2003 (04.08.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
02078231.4 2 August 2002 (02.08.2002) EP
60/427,862 20 November 2002 (20.11.2002) US
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **TIBOTEC PHARMACEUTICALS LTD.** [IE/IE]; Unit 4, Block 4B, Blanchardstown Corporate Park, Blanchardstown 15, Dublin (IE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **SURLERAUX, Dominique, Louis, Nestor, Ghislain** [BE/BE]; Diegemstraat 48, B-1830 Machelen (BE). **WIGERINCK, Piet, Tom, Bert, Paul** [BE/BE]; Kardinaal Cardijnstraat 29, B-2840 Terhagen (BE). **GETMAN, Daniel, P.** [US/US]; 66 Sunnyside Court, Chesterfield, MO 63017 (US).
- Declaration under Rule 4.17:**
— of inventorship (*Rule 4.17(iv)*) for US only
- Published:**
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **BROADSPECTRUM 2-AMINO-BENZOTHAZOLE SULFONAMIDE HIV PROTEASE INHIBITORS**



(57) **Abstract:** The present invention relates to the use of 2-amino-benzothiazoles, having the formula (I) wherein R₁ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C₁₋₆-alkyl, hydroxy, amino, halogen, aminoC₁₋₄-alkyl and mono- or di(C₁₋₄-alkyl)amino; R₂ is hydrogen or

C₁₋₆-alkyl; L is a direct bond, -O-, C₁₋₆-alkanediyl-O- or -O-C₁₋₆-alkanediyl; R₃ is phenylC₁₋₄-alkyl; R₄ is C₁₋₆-alkyl; R₅ is hydrogen or C₁₋₆-alkyl; R₆ is hydrogen or C₁₋₆-alkyl; in the manufacture of a medicament useful for inhibiting mutant HIV protease in a mammal infected with said mutant HIV protease. It also relates to novel compounds of formula (I).

BEST AVAILABLE COPY

WO 2004/014371 A1

BROADSPECTRUM 2-AMINO-BENZOTHAZOLE SULFONAMIDE HIV
PROTEASE INHIBITORS

5 This application claims priority benefit to EP Application EP 02078231.4 filed on
August 2, 2002 and to U.S. Provisional Application No. 60/427,862, filed on
November 20, 2002, the contents of which are expressly incorporated by reference
herein.

10 The present invention relates to 2-amino-benzothiazole sulfonamides, their use as
broadpectrum HIV protease inhibitors, processes for their preparation as well as
pharmaceutical compositions and diagnostic kits comprising them. The present
invention also concerns combinations of the present 2-aminobenzoxazole sulfonamides
with another anti-retroviral agent. It further relates to their use in assays as reference
compounds or as reagents.

15 The virus causing the acquired immunodeficiency syndrome (AIDS) is known by
different names, including T-lymphocyte virus III (HTLV-III) or lymphadenopathy-
associated virus (LAV) or AIDS-related virus (ARV) or human immunodeficiency
virus (HIV). Up until now, two distinct families have been identified, i.e. HIV-1 and
20 HIV-2. Hereinafter, HIV will be used to generically denote these viruses.

One of the critical pathways in a retroviral life cycle is the processing of polyprotein
precursors by aspartic protease. For instance, with the HIV virus the *gag-pol* protein is
processed by HIV protease. The correct processing of the precursor polyproteins by
25 the aspartic protease is required for the assembly of infectious virions, thus making the
aspartic protease an attractive target for antiviral therapy. In particular for HIV
treatment, the HIV protease is an attractive target.

30 HIV protease inhibitors (PIs) are commonly administered to AIDS patients in
combination with other anti-HIV compounds such as, for instance nucleoside reverse
transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors
(NNRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs) or other protease
inhibitors. Despite the fact that these antiretrovirals are very useful, they have a
common limitation, namely, the targeted enzymes in the HIV virus are able to mutate
35 in such a way that the known drugs become less effective, or even ineffective against
these mutant HIV viruses. Or, in other words, the HIV virus creates an ever-increasing
resistance against the available drugs.

-2-

Resistance of retroviruses, and in particular the HIV virus, against inhibitors is a major cause of therapy failure. For instance, half of the patients receiving anti-HIV combination therapy do not respond fully to the treatment, mainly because of resistance of the virus to one or more drugs used. Moreover, it has been shown that resistant virus is carried over to newly infected individuals, resulting in severely limited therapy options for these drug-naïve patients. On the International AIDS Conference in Paris in July 2003 researchers released that the biggest study so far of resistance to AIDS drugs finds that about 10 percent of all newly infected people in Europe have drug-resistant strains. Smaller tests to determine the spread of resistance have been done in the high-risk city center of San Francisco. This test showed the highest level of resistance at 27 percent. Therefore, there is a need in the art for new compounds for retrovirus therapy, more particularly for AIDS therapy. The need in the art is particularly acute for compounds that are active not only on wild type HIV virus, but also on the increasingly more common resistant HIV viruses.

Known antiretrovirals, often administered in a combination therapy regimen, will eventually cause resistance as stated above. This often may force the physician to boost the plasma levels of the active drugs in order for said antiretrovirals to regain effectivity against the mutated HIV viruses. The consequence of which is a highly undesirable increase in pill burden. Boosting plasma levels may also lead to an increased risk of non-compliance with the prescribed therapy. Thus, it is not only important to have compounds showing activity for a wide range of HIV mutants, it is also important that there is little or no variance in the ratio between activity against mutant HIV virus and activity against wild type HIV virus (also defined as fold resistance or FR) over a broad range of mutant HIV strains. As such, a patient may remain on the same combination therapy regimen for a longer period of time since the chance that a mutant HIV virus will be sensitive to the active ingredients will be increased.


Finding compounds with a high potency on the wild type and on a wide variety of mutants is also of importance since the pill burden can be reduced if therapeutic levels are kept to a minimum. One additional way of reducing this pill burden is finding anti-HIV compounds with good bioavailability, i.e. a favorable pharmacokinetic and metabolic profile, such that the daily dose can be minimized and consequently also the number of pills to be taken.

Another favorable characteristic of an anti-HIV compound is that plasma protein binding of the inhibitor has minimal or even no effect on its potency.


Thus, there is a high medical need for protease inhibitors that are able to combat a broad spectrum of mutants of the HIV virus with little variance in fold resistance. Those protease inhibitors with a good bioavailability and little or no effect on their potency due to plasma protein binding have additional advantages.

5

Up until now, several protease inhibitors are on the market or are being developed. One particular core structure (depicted below) has been disclosed in a number of references, such as, WO 95/06030, WO 96/22287, WO 96/28418, WO 96/28463, WO 96/28464, WO 96/28465 and WO 97/18205. The compounds disclosed therein are described as retroviral protease inhibitors.




WO 99/67254 discloses 4-substituted-phenyl sulfonamides capable of inhibiting multi-drug resistant retroviral proteases.



15 Surprisingly, the 2-amino-benzothiazole sulfonamides of the present invention are found to have a favorable virological profile. Not only are they active against wild-type HIV virus, but they also show a broadspectrum activity against various mutant HIV viruses exhibiting resistance against known protease inhibitors.

20 The present invention concerns the use of 2-amino-benzothiazole protease inhibitors,
having the formula



and *N*-oxides, salts, stereoisomeric forms, racemic mixtures, prodrugs, esters and metabolites thereof, wherein

25 R₁ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently

selected from C₁₋₆alkyl, hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono-or di(C₁₋₄alkyl)amino;

R₂ is hydrogen or C₁₋₆alkyl;

L is a direct bond, -O-, C₁₋₆alkanediyl-O- or -O-C₁₋₆alkanediyl;

5 R₃ is phenylC₁₋₄alkyl;

R₄ is C₁₋₆alkyl;

R₅ is hydrogen or C₁₋₆alkyl;

R₆ is hydrogen or C₁₋₆alkyl;

10 in the manufacture of a medicament useful for inhibiting mutant HIV protease in a mammal infected with said mutant HIV protease. Said mammal in particular is a human being. The compounds of the present invention are in particular useful in the manufacture of a medicament useful for inhibiting a broad range of mutant HIV proteases.

15 A special interest goes to the free base, salt or N-oxide form of the compounds of formula (I), and their stereoisomeric forms.

Also of special interest is the use of the present compounds wherein R₁ is tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with
20 one or more substituents independently selected from C₁₋₆alkyl, hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono-or di(C₁₋₄alkyl)amino in the manufacture of a medicament useful for inhibiting mutant HIV protease in a mammal infected with said mutant HIV protease.

25 A mutant of the HIV protease enzyme is defined as an HIV protease enzyme which has at least one mutation in its amino acid sequence relative to the amino acid sequence of the wild-type HIV protease. For purposes of denoting the mutants throughout the text, the HXB2 wild type reference (HIV IIIB LAI wild type), of which the sequence can be found at NIH's GenBank, is used.

30

The standard of "sensitivity" or alternatively "resistance" of a HIV protease enzyme to a drug is set by the commercially available HIV protease inhibitors. As explained hereinabove, existing commercial HIV protease inhibitors may loose effectivity over time against a population of HIV virus in a patient. The reason being that under
35 pressure of the presence of a particular HIV protease inhibitor, the existing population of HIV virus, often mainly wild type HIV protease enzyme, mutates into different mutants which may be less sensitive to that same HIV protease inhibitor. If this phenomenon occurs, one talks about resistant mutants. If those mutants are not only

resistant to that one particular HIV protease inhibitor, but also to one or multiple other commercially available HIV protease inhibitors, one talks about multi-drug resistant HIV protease. One way of expressing the resistance of a mutant to a particular HIV protease inhibitor is making the ratio between the EC_{50} of said HIV protease inhibitor against mutant HIV protease over EC_{50} of said HIV protease inhibitor against wild type HIV protease. Said ratio is also called fold resistance (FR).

Many of the mutants occurring in the clinic have a fold resistance of 100 or more against the commercially available HIV protease inhibitors, like saquinavir, indinavir, ritonavir and nelfinavir. Clinically relevant mutants of the HIV protease enzyme can for instance be characterized by a mutation at amino acid position 10, 71 and/or 84. Examples of such clinical relevant mutant HIV proteases are listed in Table 1.

The compounds of the present invention show a fold resistance ranging between 0.01 and 100 against at least one, often against a broad range, of clinically relevant mutant HIV proteases. A particular group of compounds of formula (I) are those compounds of formula (I) showing a fold resistance against at least one mutant HIV protease ranging between 0.1 and 100, suitably ranging between 0.1 and 50, and more suitably ranging between 0.1 and 30. Of particular interest are the compounds of formula (I) showing a fold resistance against at least one mutant HIV protease ranging between 0.1 and 20, and even more interesting are those compounds of formula (I) showing a fold resistance against at least one mutant HIV protease ranging between 0.1 and 10.

Thus, the present invention relates to the use of a compound of formula (I) in the manufacture of a medicament useful for inhibiting replication of a HIV virus having a mutant HIV protease, in particular a multi-drug resistant mutant HIV protease. It also relates to the use of a compound of formula (I) in the manufacture of a medicament useful for treating or combating a disease associated with HIV viral infection wherein the protease of the HIV virus is mutant, in particular a multi-drug resistant mutant HIV protease.

In other words, the present invention relates to a method of inhibiting a mutant HIV protease, in particular a multi-drug resistant mutant HIV protease, in a mammal infected with said mutant HIV protease, said method comprising contacting said mutant HIV protease in said mammal with an effective amount of a compound of formula (I). The present invention also relates to a method of inhibiting replication of a HIV virus, which has a mutant HIV protease, in particular a multi-drug resistant mutant HIV protease, in a mammal, said method comprising contacting said HIV virus, which has a mutant HIV protease, in said mammal with an effective amount of a compound of

formula (I). The present invention further relates to a method of treating or combating a mammalian disease associated with HIV viral infection wherein the protease of the HIV virus is mutant, in particular a multi-drug resistant mutant HIV protease, said method comprising contacting said HIV virus wherein the protease of the HIV virus is mutant infecting said mammal with an effective amount of a compound of formula (I).

Of particular interest is that the compounds of the present invention can be used in the manufacture of a medicament for the treatment of individuals infected with mutant HIV protease bearing a mutation at least at one of the amino acid positions 10, 71 or 84 or at least a combination of two of these positions or at least a combination of all three.

A basic nitrogen occurring in the present compounds can be quaternized with any agent known to those of ordinary skill in the art including, for instance, lower alkyl halides, dialkyl sulfates, long chain halides and aralkyl halides.

Whenever the term "substituted" is used in defining the compounds of formula (I), it is meant to indicate that one or more hydrogens on the atom(s) indicated in the expression using "substituted" is replaced with a selection from the indicated group, provided that the indicated atom's normal valency is not exceeded, and that the substitution results in a chemically stable compound, i.e. a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into a therapeutic agent.

As used herein, the term "halo" or "halogen" as a group or part of a group is generic for fluoro, chloro, bromo or iodo.

The term "C₁₋₄alkyl" as a group or part of a group defines straight and branched chained saturated hydrocarbon radicals having from 1 to 4 carbon atoms, such as, for example, methyl, ethyl, propyl, butyl and 2-methyl-propyl.

The term "C₁₋₆alkyl" as a group or part of a group defines straight and branched chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the groups defined for C₁₋₄alkyl and pentyl, hexyl, 2-methylbutyl, 3-methylpentyl and the like.

The term "C₁₋₆alkanediyl" as a group or part of a group defines bivalent straight and branched chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, methylene, ethan-1,2-diyl, propan-1,3-diyl, propan-1,2-diyl, butan-1,4-diyl, pentan-1,5-diyl, hexan-1,6-diyl, 2-methylbutan-1,4-diyl, 3-methyl-pentan-1,5-diyl and the like.

As used herein, the term "one or more" covers the possibility of all the available atoms, where appropriate, to be substituted, preferably, one, two or three.

The term "prodrug" as used throughout this text means the pharmacologically acceptable derivatives such as esters, amides and phosphates, such that the resulting *in vivo* biotransformation product of the derivative is the active drug as defined in the compounds of formula (I). The reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8th ed, McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p 13–15) describing prodrugs generally is hereby incorporated. Prodrugs of a compound of the present invention are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of the present invention wherein a hydroxy group, for instance the hydroxy group on the asymmetric carbon atom, or an amino group is bonded to any group that, when the prodrug is administered to a patient, cleaves to form a free hydroxyl or free amino, respectively.

Typical examples of prodrugs are described for instance in WO 99/33795, WO 99/33815, WO 99/33793 and WO 99/33792 all incorporated herein by reference. Prodrugs are typically characterized by excellent aqueous solubility, increased bioavailability and are readily metabolized into the active inhibitors *in vivo*.

For therapeutic use, the salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically or physiologically acceptable. However, salts having a pharmaceutically unacceptable counterion may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound of formula (I). All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

The pharmaceutically acceptable or physiologically tolerable addition salt forms which the compounds of the present invention are able to form can conveniently be prepared using the appropriate acids, such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; hemisulphuric, nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, aspartic, dodecylsulphuric, heptanoic, hexanoic, nicotinic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methane-sulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

Conversely said acid addition salt forms can be converted by treatment with an appropriate base into the free base form.

- 5 The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt form by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, N-methyl, -D-glucamine, hydrabamine salts, and salts with amino acids
10 such as, for example, arginine, lysine and the like.

Conversely said base addition salt forms can be converted by treatment with an appropriate acid into the free acid form.

15

The term "salts" also comprises the hydrates and the solvent addition forms that the compounds of the present invention are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

- 20 The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

The present compounds may also exist in their tautomeric forms. Such forms, although not explicitly indicated in the above formula are intended to be included within the
25 scope of the present invention.


- The term stereochemically isomeric forms of compounds of the present invention, as used hereinbefore, defines all possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures which
30 are not interchangeable, which the compounds of the present invention may possess. Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of all possible stereochemically isomeric forms that said compound might possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically
35 isomeric forms of the compounds of the present invention both in pure form and in admixture with each other are intended to be embraced within the scope of the present invention.

Pure stereoisomeric forms of the compounds and intermediates as mentioned herein are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure of said compounds or intermediates. In particular, the term 'stereoisomerically pure' concerns compounds or intermediates having a stereoisomeric excess of at least 80% (i. e. minimum 90% of one isomer and maximum 10% of the other possible isomers) up to a stereoisomeric excess of 100% (i.e. 100% of one isomer and none of the other), more in particular, compounds or intermediates having a stereoisomeric excess of 90% up to 100%, even more in particular having a stereoisomeric excess of 94% up to 100% and most in particular having a stereoisomeric excess of 97% up to 100%. The terms 'enantiomerically pure' and 'diastereomerically pure' should be understood in a similar way, but then having regard to the enantiomeric excess, respectively the diastereomeric excess of the mixture in question.

Pure stereoisomeric forms of the compounds and intermediates of this invention may be obtained by the application of art-known procedures. For instance, enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids or bases. Examples thereof are tartaric acid, dibenzoyltartaric acid, ditoluoyltartaric acid and camphosulfonic acid. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably, if a specific stereoisomer is desired, said compound would be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

The diastereomeric racemates of formula (I) can be obtained separately by conventional methods. Appropriate physical separation methods that may advantageously be employed are, for example, selective crystallization and chromatography, e.g. column chromatography.

It is clear to a person skilled in the art that the compounds of formula (I) contain at least two asymmetric centers and thus may exist as different stereoisomeric forms. These two asymmetric centers are indicated with an asterisk (*) in the figure below.



The absolute configuration of each asymmetric center that may be present in the compounds of formula (I) may be indicated by the stereochemical descriptors R and S, this R and S notation corresponding to the rules described in Pure Appl. Chem. 1976, 45, 11-30. The carbon atom bearing the hydroxy group and marked with the asterisk (*) preferably has the R configuration. The carbon atom bearing the R³ group and marked with the asterisk (*) preferably has the S configuration.

The present invention is also intended to include all isotopes of atoms occurring on the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

Whenever used hereinafter, the term "compounds of formula (I)", or "the present compounds" or similar term is meant to include the compounds of general formula (I), their *N*-oxides, salts, stereoisomeric forms, racemic mixtures, prodrugs, esters and metabolites, as well as their quaternized nitrogen analogues.

Some of the compounds of formula (I) have been disclosed in WO 95/06030, i.e.

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid benzyl ester;

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid pyridin-3-ylmethyl ester;

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid thiazol-5-ylmethyl ester;

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;

3-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;

4-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;

5-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;

N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;


N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-4-hydroxy-2-methyl-benzamide;

N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-3-hydroxy-2-methyl-benzamide; and

-11-

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (S)-(tetrahydrofuran-3-yl) ester.

Hence, the present invention also concerns the compounds of formula (I)



and *N*-oxides, salts, stereoisomeric forms, racemic mixtures, prodrugs, esters and metabolites thereof, wherein

R_1 is hexahydrofuro[2,3-*b*]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C_{1-6} alkyl, hydroxy, amino, halogen, amino C_{1-4} alkyl and mono- or di(C_{1-4} alkyl)amino;

R_2 is hydrogen or C_{1-6} alkyl;

L is a direct bond, -O-, C_{1-6} alkanediyl-O- or -O- C_{1-6} alkanediyl;

R_3 is phenyl C_{1-4} alkyl;

R_4 is C_{1-6} alkyl;

R_5 is hydrogen or C_{1-6} alkyl;

R_6 is hydrogen or C_{1-6} alkyl;

provided that the compounds are other than :

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid benzyl ester;

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid pyridin-3-ylmethyl ester;

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid thiazol-5-ylmethyl ester;

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;

3-amino- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;

4-amino- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;

5-amino- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;

N- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;

- N-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-4-hydroxy-2-methyl-benzamide;
N-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-hydroxy-2-methyl-benzamide; and
5 {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (S)-(tetrahydrofuran-3-yl) ester.

10 Interesting compounds are those compounds of formula (I) wherein R¹ is hexahydrofuro[2,3-b]furanyl or oxazolyl.

Other interesting compounds are those compounds of formula (I) or those compounds belonging to any subgroup thereof wherein R₁ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, and L is a direct bond.

- 15 Yet other interesting compounds are those compounds of formula (I) or those compounds belonging to any subgroup thereof wherein R₁ is hexahydrofuro[2,3-b]furanyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C₁₋₆alkyl, hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono-or di(C₁₋₄alkyl)amino; and L is -O-.

- 20 Still other interesting compounds are those compounds of formula (I) or those compounds belonging to any subgroup thereof wherein R₁ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, or phenyl substituted with one or more substituents independently selected from C₁₋₆alkyl, hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono-or di(C₁₋₄alkyl)amino; and L is C₁₋₆alkanediyl-O- whereby the -O- is
25 attached to the nitrogen of the amide.

- Also interesting compounds are those compounds of formula (I) or those compounds belonging to any subgroup thereof wherein R₁ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with
30 one or more substituents independently selected from hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono-or di(C₁₋₄alkyl)amino; and L is -O-C₁₋₆alkanediyl whereby -O- is attached to the R¹ group.

- 35 A suitable group of compounds are those compounds of formula (I) or those compounds belonging to any subgroup thereof wherein at least one of R₅ and R₆ is C₁₋₆alkyl, in particular R₅ is methyl and R₆ is hydrogen or methyl, more in particular R₅ is methyl and R₆ is hydrogen.

Compounds of particular interest are those compounds of formula (I) or those compounds belonging to any subgroup thereof wherein $-L-R^1$ is $-O-(\text{hexahydrofuro}[2,3-b]\text{furan})$, $-O-\text{tetrahydrofuran}$, $-O-\text{methyl}-(\text{optionally substituted phenyl})$, $-O-\text{methyl-pyridinyl}$, $-O-\text{methyl-thiazolyl}$, $-O-\text{methyl-oxazolyl}$, $-\text{methyl-O}-(\text{optionally substituted phenyl})$ or optionally substituted phenyl. Preferably, the optional substituents on the phenyl group are methyl, amino, hydroxy, halogen, aminomethyl,

Compounds of special interest are those compounds of formula (I) or those compounds belonging to any subgroup thereof wherein R^1 is hexahydrofuro[2,3-b]furan, tetrahydrofuran, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from $C_{1-6}\text{alkyl}$, hydroxy, amino, chloro, bromo, amino $C_{1-4}\text{alkyl}$ and mono- or di($C_{1-4}\text{alkyl}$)amino.

Suitably, one or more of the following restrictions apply to any of the above mentioned interesting subgroups of the compounds of formula (I) or subgroups of particular or special interest:

R^2 is hydrogen;

R^3 is phenylmethyl;

R^4 is $C_{1-4}\text{alkyl}$, preferably isobutyl;

R^5 is hydrogen or methyl;

R^6 is hydrogen or methyl.

An interesting combination for a compound of formula (I) or a compound of any subgroup thereof is formed by R^2 being hydrogen; R^3 being phenylmethyl and R^4 being $C_{1-4}\text{alkyl}$, preferably isobutyl;

A special subgroup of the compounds of formula (I) is defined as encompassing those compounds of formula (I) wherein R^5 and R^6 are both hydrogen.

Another special subgroup of the compounds of formula (I) or of the compounds belonging to any subgroup thereof are those compounds wherein $-L-R^1$ is $-O-(\text{hexahydrofuro}[2,3-b]\text{furan})$, $-O-\text{tetrahydrofuran}$, $-O-\text{methyl-thiazolyl}$, $-O-\text{methyl-oxazolyl}$, $-\text{methyl-O}-(2,6-\text{dimethylphenyl})$, $-\text{methyl-O}-(4-\text{aminomethyl-2,6-dimethylphenyl})$, $-\text{methyl-O}-(4-\text{amino-2,6-dimethylphenyl})$, 3-hydroxy-2-methyl-phenyl or 3-amino-2-methyl-phenyl; and R^5 is methyl or hydrogen and R^6 is hydrogen.

Preferred compounds are

- {3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid thiazol-5-ylmethyl ester;
- 5 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
- 10 carbamic acid benzyl ester;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid pyridin-3-ylmethyl ester;
- 15 3-amino-N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-hydroxy-2-methyl-benzamide;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
- 20 carbamic acid tetrahydro-furan-3-yl ester;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
- 25 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-fluoro-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-aminomethyl-2,6-dimethyl-phenoxy)-acetamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
- 30 propyl}-carbamic acid thiazol-5-ylmethyl ester;
3-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-2-methyl-benzamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid tetrahydro-furan-3-yl ester;
- 35 N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-iodo-2,6-dimethyl-phenoxy)-acetamide;

- 2-(4-aminomethyl-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
2-(4-amino-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
5 N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-4-bromo-2-methyl-benzamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid oxazol-5-ylmethyl ester;
4-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide; their *N*-oxides and salts and
10 the stereoisomeric forms thereof.

- Another group of interest includes
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
15 {1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
20 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-fluoro-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-aminomethyl-2,6-dimethyl-phenoxy)-acetamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid thiazol-5-ylmethyl ester;
25 3-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-2-methyl-benzamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid tetrahydro-furan-3-yl ester;
30 N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-iodo-2,6-dimethyl-phenoxy)-acetamide;
2-(4-aminomethyl-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
35 2-(4-amino-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;

-16-


N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-4-bromo-2-methyl-benzamide;

{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid oxazol-5-ylmethyl ester;

- 5 4-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide; their *N*-oxides and salts and the stereoisomeric forms thereof.

- 10 Most preferred compounds are those enantiomeric forms of the compounds of formula (I) or of the compounds belonging to any subgroup thereof having a (1*S*,2*R*)-1-benzyl-2-hydroxy-propyl configuration.

- Those compounds of formula (I) or those compounds belonging to any subgroup thereof in a hexahydro-furo[2,3-*b*]furan-3-yl ester form of the carbamic acid derivative occur
15 preferably in a (3*R*,3*aS*,6*aR*) form such as, for instance, {(1*S*,2*R*)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid




(3*R*,3*aS*,6*aR*)-hexahydro-furo[2,3-*b*]furan-3-yl ester

- 20 The compounds of formula (I) can generally be prepared using procedures analogous to those procedures described in WO 95/06030, WO 96/22287, WO 96/28418, WO 96/28463, WO 96/28464, WO 96/28465 and WO 97/18205.

- 25 Particular reaction procedures to make the present compounds are described below. In the preparations described below, the reaction products may be isolated from the medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, trituration and chromatography.

Scheme A

-17-



The 2-amino-6-chlorosulfonylbenzothiazole derivative (intermediate **a-2**) was prepared following the procedure described in EP-A-0,445,926. Intermediates **a-4** were prepared by reacting an intermediate **a-3**, prepared according to the procedure described in WO97/18205 and also depicted in scheme B, with an intermediate **a-2** in a reaction-inert solvent such as dichloromethane, and in the presence of a base such as triethylamine and at low temperature, for example at 0°C. The Boc group in the intermediate **a-3** is a protective *tert*-butoxycarbonyl group. Another suitable protective group such as phthalimido or benzyloxycarbonyl may conveniently replace it.


Intermediates **a-4** may be deprotected with an acid such as hydrochloric acid in isopropanol or with trifluoroacetic acid depending on the nature of the amino group in the 2 position of benzoxazole, in a suitable solvent such as a mixture of ethanol and dioxane, thus preparing an intermediate **a-5**. Said intermediate **a-5** may be further reacted with an intermediate of formula $\text{R}_1\text{-L-C(=O)-OH}$ in the presence of a base such as triethylamine (for alcohols to generate a carbamate) and optionally in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloric acid (EDC) and 1-hydroxybenzotriazole (HOBT) (for carboxylic acids to generate an amide) or an alcohol such as *tert*-butanol, and in a suitable solvent such as dichloromethane; thus forming intermediates **a-6**.

A convenient way of preparing compounds of formula (I) wherein both R_5 and R_6 are hydrogen can be prepared analogously to the procedure described in scheme A, and whereby one of R_5 or R_6 is replaced by a suitable protective group such as, for example, an acetyl or an alkyloxycarbonyl group. In such a case, deprotection may

occur simultaneously with the deprotection of the nitrogen atom on the left-hand side of the molecule.

A number of intermediates and starting materials used in the foregoing preparations are known compounds, while others may be prepared according to art-known methodologies of preparing said or similar compounds.

Scheme B



Intermediate **b-2**, corresponding to intermediate **a-3** in scheme A, may be prepared by adding an amine of formula H_2N-R_4 to an intermediate **b-1** in a suitable solvent such as isopropanol.

In scheme B, enantiomerically pure compounds of formula **b-2** are only obtained if **b-1** is enantiomerically pure. If **b-1** is a mixture of stereoisomers, than **b-2** will also consist of a mixture of stereoisomers.

The compounds of formula (I) may also be converted to the corresponding *N*-oxide forms following art-known procedures for converting trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chloro-benzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *tert*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

The present compounds can thus be used in animals, preferably in mammals, and in particular in humans as pharmaceuticals per se, in mixtures with one another or in the form of pharmaceutical preparations.

Furthermore, the present invention relates to pharmaceutical preparations that as active constituents contain an effective dose of at least one of the compounds of formula (I) in addition to customary pharmaceutically innocuous excipients and auxiliaries. The pharmaceutical preparations normally contain 0.1 to 90% by weight of a compound of formula (I). The pharmaceutical preparations can be prepared in a manner known per se to one of skill in the art. For this purpose, at least one of a compound of formula (I), together with one or more solid or liquid pharmaceutical excipients and/or auxiliaries and, if desired, in combination with other pharmaceutical active compounds, are brought into a suitable administration form or dosage form which can then be used as a pharmaceutical in human medicine or veterinary medicine.

Pharmaceuticals which contain a compound according to the invention can be administered orally, parenterally, e.g., intravenously, rectally, by inhalation, or topically, the preferred administration being dependent on the individual case, e.g., the particular course of the disorder to be treated. Oral administration is preferred.

The person skilled in the art is familiar on the basis of his expert knowledge with the auxiliaries that are suitable for the desired pharmaceutical formulation. Beside solvents, gel-forming agents, suppository bases, tablet auxiliaries and other active compound carriers, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, agents for achieving a depot effect, buffer substances or colorants are also useful.

Due to their favorable pharmacological properties, particularly their activity against multi-drug resistant HIV protease enzymes, the compounds of the present invention are useful in the treatment of individuals infected by HIV and for the prophylaxis of these individuals. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses whose existence is mediated by, or depends upon, the protease enzyme. Conditions which may be prevented or treated with the compounds of the present invention, especially conditions associated with HIV and other pathogenic retroviruses, include AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic CNS diseases caused by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis.

Said method of treatment comprises the systemic administration to HIV-infected subjects of an amount effective to combat the conditions associated with HIV virus with multi-drug resistant protease enzyme.

The compounds of the present invention may also find use in inhibiting *ex vivo* samples containing multi-drug resistant HIV-protease or expected to be exposed to multi-drug resistant HIV-protease. Hence, the present compounds may be used to inhibit multi-
5 drug resistant HIV-protease present in a body fluid sample that contains or is suspected to contain or be exposed to multi-drug resistant HIV-protease.

Also, the combination of an antiretroviral compound and a compound of the present invention can be used as a medicine. Thus, the present invention also relates to a
10 product containing (a) a compound of the present invention, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in treatment of retroviral infections, in particular, in the treatment of infections with multi-drug resistant HIV proteases. Thus, to combat or treat infections with multi-drug resistant HIV protease, or the infection and disease associated with
15 such infections, such as Acquired Immunodeficiency Syndrome (AIDS) or AIDS Related Complex (ARC), the compounds of this invention may be co-administered in combination with for instance, binding inhibitors, such as, for example, dextran sulfate, suramine, polyanions, soluble CD4, PRO-542, BMS-806; fusion inhibitors, such as, for example, T20, T1249, 5-helix, D-peptide ADS-J1; co-receptor binding inhibitors, such
20 as, for example, AMD 3100, AMD-3465, AMD7049, AMD3451 (Bicyclams), TAK 779; SHC-C (SCH351125), SHC-D, PRO-140RT inhibitors, such as, for example, foscarnet and prodrugs; nucleoside RTIs, such as, for example, AZT, 3TC, DDC, DDI, D4T, Abacavir, FTC, DAPD, dOTC, DPC 817; nucleotide RTIs, such as, for example, PMEA, PMPA (tenofovir); NNRTIs, such as, for example, nevirapine, delavirdine,
25 efavirenz, 8 and 9-Cl TIBO (tivicar), loviride, TMC-125, dapivirine, MKC-442, UC 781, UC 782, Capravirine, DPC 961, DPC963, DPC082, DPC083, calanolide A, SJ-1366, TSAO, 4"-deaminated TSAO, MV150, MV026048; RNase H inhibitors, such as, for example, SP1093V, PD126338; TAT inhibitors, such as, for example, RO-5-3335, K12, K37; integrase inhibitors, such as, for example, L 708906, L 731988, S-
30 1360; protease inhibitors, such as, for example, amprenavir and prodrug GW908, ritonavir, nelfinavir, saquinavir, indinavir, lopinavir, palinavir, BMS 186316, atazanavir, DPC 681, DPC 684, tipranavir, AG1776, mozenavir, GS3333, KNI-413, KNI-272, L754394, L756425, LG-71350, PD161374, PD173606, PD177298, PD178390, PD178392, PNU 140135, TMC114 maslinic acid, U-140690; glycosylation
35 inhibitors, such as, for example, castanospermine, deoxynojirimycin.

The combination may in some cases provide a synergistic effect, whereby viral infectivity and its associated symptoms may be prevented, substantially reduced, or eliminated completely.

5 The compounds of the present invention may also be administered in combination with immunomodulators (e.g., bropiramine, anti-human alpha interferon antibody, IL-2, methionine enkephalin, interferon alpha, HE-2000 and naltrexone) with antibiotics (e.g., pentamidine isothiorate) cytokines (e.g. Th2), modulators of cytokines, chemokines or the receptors thereof (e.g. CCR5) or hormones (e.g. growth hormone) to
10 ameliorate, combat, or eliminate HIV infection and its symptoms. Such combination therapy in different formulations may be administered simultaneously, separately or sequentially. Alternatively, such combination may be administered as a single formulation, whereby the active ingredients are released from the formulation simultaneously or separately.

15 The compounds of the present invention may also be administered in combination with modulators of the metabolism following application of the drug to an individual. These modulators include compounds that interfere with the metabolism at cytochromes, such as cytochrome P450. Some modulators inhibit cytochrome P450. It
20 is known that several isoenzymes exist of cytochrome P450, one of which is cytochrome P450 3A4. Ritonavir is an example of a modulator of metabolism via cytochrome P450. Such combination therapy in different formulations may be administered simultaneously, separately or sequentially. Alternatively, such
25 combination may be administered as a single formulation, whereby the active ingredients are released from the formulation simultaneously or separately. Such modulator may be administered at the same or different ratio as the compound of the present invention. Preferably, the weight ratio of such modulator vis-à-vis the compound of the present invention (modulator:compound of the present invention) is 1:1 or lower, more preferable the ratio is 1:3 or lower, suitably the ratio is 1:10 or
30 lower, more suitably the ratio is 1:30 or lower.

For an oral administration form, compounds of the present invention are mixed with suitable additives, such as excipients, stabilizers or inert diluents, and brought by means of the customary methods into the suitable administration forms, such as tablets, coated
35 tablets, hard capsules, aqueous, alcoholic, or oily solutions. Examples of suitable inert carriers are gum arabic, magnesia, magnesium carbonate, potassium phosphate, lactose, glucose, or starch, in particular, cornstarch. In this case the preparation can be carried out both as dry and as moist granules. Suitable oily excipients or solvents are vegetable

or animal oils, such as sunflower oil or cod liver oil. Suitable solvents for aqueous or alcoholic solutions are water, ethanol, sugar solutions, or mixtures thereof. Polyethylene glycols and polypropylene glycols are also useful as further auxiliaries for other administration forms.

5

For subcutaneous or intravenous administration, the active compounds, if desired with the substances customary therefore such as solubilizers, emulsifiers or further auxiliaries, are brought into solution, suspension, or emulsion. The compounds of formula (I) can also be lyophilized and the lyophilizates obtained used, for example, for the production of injection or infusion preparations. Suitable solvents are, for example, water, physiological saline solution or alcohols, e.g. ethanol, propanol, glycerol, in addition also sugar solutions such as glucose or mannitol solutions, or alternatively mixtures of the various solvents mentioned.

10

15 Suitable pharmaceutical formulations for administration in the form of aerosols or sprays are, for example, solutions, suspensions or emulsions of the compounds of formula (I) or their physiologically tolerable salts in a pharmaceutically acceptable solvent, such as ethanol or water, or a mixture of such solvents. If required, the formulation can also additionally contain other pharmaceutical auxiliaries such as surfactants, emulsifiers and stabilizers as well as a propellant. Such a preparation customarily contains the active compound in a concentration from approximately 0.1 to 50%, in particular from approximately 0.3 to 3% by weight.

20

In order to enhance the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions, it can be advantageous to employ α -, β - or γ -cyclodextrins or their derivatives. Also co-solvents such as alcohols may improve the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions. In the preparation of aqueous compositions, addition salts of the subject compounds are obviously more suitable due to their increased water solubility.

25

30

Appropriate cyclodextrins are α -, β - or γ -cyclodextrins (CDs) or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C_{1-6} alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxy C_{1-6} alkyl, particularly hydroxyethyl, hydroxypropyl or hydroxybutyl; carboxy C_{1-6} alkyl, particularly carboxymethyl or carboxyethyl; C_{1-6} alkyl-carbonyl, particularly acetyl; C_{1-6} alkyloxycarbonyl C_{1-6} alkyl or carboxy C_{1-6} alkyloxy C_{1-6} alkyl, particularly carboxymethoxypropyl or carboxyethoxypropyl; C_{1-6} alkylcarbonyloxy C_{1-6} alkyl, particularly 2-acetyloxypropyl. Especially noteworthy as complexants and/or solubilizers are β -CD, randomly methylated β -CD,

35

2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- γ -CD, 2-hydroxy-propyl- γ -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxy-propyl- β -CD (2-HP- β -CD).

- 5 The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

10 An interesting way of formulating the present compounds in combination with a cyclodextrin or a derivative thereof has been described in EP-A-721,331. Although the formulations described therein are with antifungal active ingredients, they are equally interesting for formulating the compounds of the present invention. The formulations described therein are particularly suitable for oral administration and comprise an antifungal as active ingredient, a sufficient amount of a cyclodextrin or a derivative
15 thereof as a solubilizer, an aqueous acidic medium as bulk liquid carrier and an alcoholic co-solvent that greatly simplifies the preparation of the composition. Said formulations may also be rendered more palatable by adding pharmaceutically acceptable sweeteners and/or flavors.

- 20 Other convenient ways to enhance the solubility of the compounds of the present invention in pharmaceutical compositions are described in WO 94/05263, WO 98/42318, EP-A-499,299 and WO 97/44014, all incorporated herein by reference.

25 More in particular, the present compounds may be formulated in a pharmaceutical composition comprising a therapeutically effective amount of particles consisting of a solid dispersion comprising (a) a compound of formula (I), and (b) one or more pharmaceutically acceptable water-soluble polymers.

- 30 The term "a solid dispersion" defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, wherein one component is dispersed more or less evenly throughout the other component or components. When said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermodynamics, such a solid dispersion is referred to as "a solid solution". Solid solutions are
35 preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered.

The term "a solid dispersion" also comprises dispersions that are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically

uniform throughout or comprise more than one phase.

The water-soluble polymer in the particles is conveniently a polymer that has an apparent viscosity of 1 to 100 mPa.s when dissolved in a 2 % aqueous solution at 20°C solution.

Preferred water-soluble polymers are hydroxypropyl methylcelluloses or HPMC. HPMC having a methoxy degree of substitution from about 0.8 to about 2.5 and a hydroxypropyl molar substitution from about 0.05 to about 3.0 is generally water soluble. Methoxy degree of substitution refers to the average number of methyl ether groups present per anhydroglucose unit of the cellulose molecule. Hydroxy-propyl molar substitution refers to the average number of moles of propylene oxide that have reacted with each anhydroglucose unit of the cellulose molecule.

First preparing a solid dispersion of the components, and then optionally grinding or milling that dispersion allows one to prepare the particles as defined hereinabove. Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation, melt-extrusion being preferred.

It may further be convenient to formulate the present compounds in the form of nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm. Useful surface modifiers are believed to include those that physically adhere to the surface of the antiretroviral agent but do not chemically bond to the antiretroviral agent.

Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

Yet another interesting way of formulating the present compounds involves a pharmaceutical composition whereby the present compounds are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition with good bioavailability which can conveniently be manufactured and which is suitable for preparing pharmaceutical dosage forms for oral administration.

Said beads comprise (a) a central, rounded or spherical core, (b) a coating film of a hydrophilic polymer and an antiretroviral agent and (c) a seal-coating polymer layer.

Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.

5

The route of administration may depend on the condition of the subject, co-medication and the like.

Another aspect of the present invention concerns a kit or container comprising a compound of formula (I) in an amount effective for use as a standard or reagent in a
10 test or assay for determining the ability of a potential pharmaceutical to inhibit multi-drug resistant HIV protease, HIV growth, or both. This aspect of the invention may find its use in pharmaceutical research programs.

The compounds of the present invention can be used in phenotypic resistance
15 monitoring assays, such as known recombinant assays, in the clinical management of resistance developing diseases such as HIV. A particularly useful resistance monitoring system is a recombinant assay known as the AntivirogramTM. The AntivirogramTM is a highly automated, high throughput, second generation, recombinant assay that can measure susceptibility, especially viral susceptibility, to the
20 compounds of the present invention. (Hertogs K, de Bethune MP, Miller V *et al.* *Antimicrob Agents Chemother*, 1998; 42(2):269-276, incorporated by reference).

Interestingly, the compounds of the present invention may comprise chemically reactive moieties capable of forming covalent bonds to localized sites such that said
25 compound have increased tissue retention and half-lives. The term "chemically reactive group" as used herein refers to chemical groups capable of forming a covalent bond. Reactive groups will generally be stable in an aqueous environment and will usually be carboxy, phosphoryl, or convenient acyl group, either as an ester or a mixed anhydride, or an imidate, or a maleimide thereby capable of forming a covalent bond with
30 functionalities such as an amino group, a hydroxy or a thiol at the target site on for example blood components such as albumine. The compounds of the present invention may be linked to maleimide or derivatives thereof to form conjugates.

The dose of the present compounds or of the physiologically tolerable salt(s) thereof to
35 be administered depends on the individual case and, as customary, is to be adapted to the conditions of the individual case for an optimum effect. Thus it depends, of course, on the frequency of administration and on the potency and duration of action of the compounds employed in each case for therapy or prophylaxis, but also on the nature

and severity of the infection and symptoms, and on the sex, age, weight, co-medication, and individual responsiveness of the human or animal to be treated and on whether the therapy is acute or prophylactic. Customarily, the daily dose of a compound of formula (I) in the case of administration to a patient approximately 75 kg in weight is 1 mg to 3g, suitably 1 mg to 1g, preferably 3 mg to 0.5 g, more preferably 5 mg to 300 mg. The dose can be administered in the form of an individual dose, or divided into several, e.g. two, three, or four, individual doses.

Experimental Part

10 Preparation of the compounds of formula (I)

The nomenclature used throughout the description is based on Chemical Abstracts Services Nomenclature.

Example 1 : Compound 2

15 To a mixture of 825 mg 2-amino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide and 373 mg triethylamine in dichloromethane was added 452 mg 1-[[[(3R,3aS,6aR)-hexahydrofuro[2,3-b]furan-3-yl]oxy]carbonyl]oxy]-2,5-pyrrolidine-dione (described in W09967417). This mixture was stirred at room temperature for 12 hours. After evaporation of dichloromethane under reduced pressure, 20 the crude product was purified on silica, yielding 270 mg 24.8 % compound 2.

Example 2 : Compound 4

To a mixture of 350 mg 2-methylamino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide and 200 mg triethylamine in dichloromethane 25 was added 210 mg 1-[[[hexahydrofuro[2,3-b]furan-3-yl]oxy]carbonyl]oxy]-2,5-pyrrolidinedione (described in W09967417). This mixture was stirred at room temperature for 12 hours. After evaporation of dichloromethane under reduced pressure, the crude product was purified on silica, yielding 260 mg (55 %) of compound 4.

30 Example 3 : Compound 6

To a mixture of 420 mg 2-dimethylamino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide and 98 mg triethylamine in dichloromethane was added 230 mg 1-[[[hexahydrofuro[2,3-b]furan-3-yl]oxy]carbonyl]oxy]-2,5-pyrrolidinedione (described in W09967417). This mixture was stirred at room 35 temperature for 12 hours. After evaporation of dichloromethane under reduced pressure, the crude product was purified on silica, yielding 500 mg 90 % of compound 6.

Example 4 : Compound 17

-27-

A mixture of 800 mg of 2-amino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide, 50 mg of HOBT (hydroxybenzotriazol), 420 mg of EDC and 668 mg of (3,4,5-trimethyl-benzyl)-carbamic acid tert-butyl ester compound with hydroxy acetic acid in 80 ml of dichloromethane, was stirred overnight at room temperature. The reaction mixture was then washed with water and brine. The organic layer was separated, dried and the solvent evaporated. The residue was purified by column chromatography, yielding 1 g (75 %) of [4-({3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropylcarbamoyl}-methoxy)-3,5-dimethyl-benzyl]-carbamic acid tert-butyl ester. This intermediate (500 mg) was further dissolved in methanol (20 ml) and 10 ml of a solution of HCl in isopropanol (5 to 6 N) was added dropwise. The mixture was stirred overnight at room temperature. The solvent was evaporated and the residue was purified on silica yielding 190 mg of compound 17 (43%).

15 Example 5 : Compound 27

A mixture of 134 mg of 2-methylamino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide, 4 mg of HOBT (hydroxybenzotriazol), 66 mg of EDC and 63 mg of 4-bromo-2-methyl benzoic acid in dichloromethane, was stirred overnight at room temperature. The reaction mixture was then washed with water and brine. The organic layer was separated, dried and the solvent evaporated. The residue was purified by preparative HPLC yielding 25 mg (13 %) of compound 27.

Example 6 : Compound 28

To a mixture of 4.48 g 2-methylamino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide and 2.73 g triethylamine in dichloromethane was added 3.45 g carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester oxazol-5-ylmethyl ester. This mixture was stirred at room temperature for 12 hours. After evaporation of dichloromethane under reduced pressure, the crude product was purified on silica, yielding 1.02 g 19 % compound 28.

The compounds in Table 1, not intended to be limiting the scope of the present invention, have been prepared analogous to one of the above examples and tested in support of the presently claimed invention :

Table 1

Name	Number
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester	1
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-	2

hydroxypropyl}-carbamic acid (3R,3aS,6aR)-hexahydro-furo[2,3-b]furan-3-yl ester	
{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid thiazol-5-ylmethyl ester	3
{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester	4
{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid (3R,3aS,6aR)-(hexahydro-furo[2,3-b]furan-3-yl) ester	5
{{(1S,2R)-1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxy-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester	6
{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid benzyl ester	7
N-{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide	8
{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid pyridin-3-ylmethyl ester	9
3-amino-N-{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide	10
N-{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-hydroxy-2-methyl-benzamide	11
{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid tetrahydro-furan-3-yl ester	12
{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (S) (tetrahydro-furan-3-yl) ester	13
N-{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide	14
N-{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-2-(2,6-dimethyl-phenoxy)-acetamide;	15
N-{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-fluoro-2-methyl-benzamide	16
N-{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-aminomethyl-2,6-dimethyl-phenoxy)-acetamide	17
{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid thiazol-5-ylmethyl ester	18
{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid thiazol-5-ylmethyl ester trifluoroacetate	19
3-amino-N-{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-	20

benzothiazole-6-sulfonyl)-amino]-propyl}-2-methyl-benzamide trifluoroacetate	
{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid (S) (tetrahydro-furan-3-yl) ester	21
{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid (S) (tetrahydro-furan-3-yl) ester trifluoroacetate	22
N-{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide trifluoroacetate	23
N-{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-iodo-2,6-dimethyl-phenoxy)-acetamide	24
2-(4-aminomethyl-2,6-dimethyl-phenoxy)-N-{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide	25
2-(4-amino-2,6-dimethyl-phenoxy)-N-{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide	26
N-{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-4-bromo-2-methyl-benzamide	27
{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid oxazol-5-ylmethyl ester	28
4-amino-N-{{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide	29

Antiviral analyses:

- The compounds of the present invention were examined for anti-viral activity in a cellular assay. The assay demonstrated that these compounds exhibited potent anti-HIV activity against a wild type laboratory HIV strain (HIV-1 strain LAI). The cellular assay was performed according to the following procedure.

Cellular Assay Experimental Method:

- HIV- or mock-infected MT4 cells were incubated for five days in the presence of various concentrations of the inhibitor. At the end of the incubation period, the replicating virus in the control cultures has killed all HIV-infected cells in the absence of any inhibitor. Cell viability is measured by measuring the concentration of MTT, a yellow, water soluble tetrazolium dye that is converted to a purple, water insoluble formazan in the mitochondria of living cells only. Upon solubilization of the resulting formazan crystals with isopropanol, the absorbance of the solution is monitored at 540nm. The values correlate directly to the number of living cells remaining in the culture at the completion of the five-day incubation. The inhibitory activity of the compound was monitored on the virus-infected cells and was expressed as EC₅₀ and

EC₉₀. These values represent the amount of the compound required to protect 50% and 90%, respectively, of the cells from the cytopathogenic effect of the virus. The toxicity of the compound was measured on the mock-infected cells and was expressed as CC₅₀, which represents the concentration of compound required to inhibit the growth of the cells by 50%. The selectivity index (SI) (ratio CC₅₀/EC₅₀) is an indication of the selectivity of the anti-HIV activity of the inhibitor. Wherever results are reported as e.g. pEC₅₀ or pCC₅₀ values, the result is expressed as the negative logarithm of the result expressed as EC₅₀ or CC₅₀ respectively.

10. Antiviral spectrum:

Because of the increasing emergence of drug resistant HIV strains, the present compounds were tested for their potency against clinically isolated HIV strains harboring several mutations (Table 2 and 3). These mutations are associated with resistance to protease inhibitors and result in viruses that show various degrees of phenotypic cross-resistance to the currently commercially available drugs such as for instance saquinavir, ritonavir, nelfinavir, indinavir and amprenavir.

Table 2 List with a representative selection of mutant HIV strains (A to F).

Strain	Mutations in HIV protease gene
A	V003I, L010I, V032T, L033M, E035D, S037Y, S037D, M046I, R057R/K, Q058E, L063P, K070T, A071V, I072V, I084V, L089V
B	V003I, L010I, K020R, E035D, M036I, S037N, Q058E, I062V, L063P, A071V, I072M, G073S, V077I, I084V, I085V, L090M
C	V003I, L010I, I015V, L019I, K020M, S037N, R041K, I054V, Q058E, L063P, A071V, I084V, L090M, I093L
D	V003I, L010L/I, I013V, L033I, E035D, M036I, M046L, K055R, R057K, L063P, I066F, A071V, I084V, N088D, L090M
E	V003I, L010I, V011I, A022V, L024I, E035D, M036I, S037T, R041K, I054V, I062V, L063P, A071V, I084V
F	L010F, M046I, M071V, I084V

Results:

As a measure of the broad spectrum activity of the present compounds, Table 3 shows the results of the antiviral testing in terms of pEC₅₀. (= -log of EC₅₀). The fold resistance (FR), defined as FR = EC₅₀(mutant strain)/EC₅₀(HIV-1 strain LAI) is listed in Table 4. For most of the compounds the fold resistance ranges between 0.1 and 100. Thus, the present compounds are potent inhibitors of a broad range of mutant strains.

-31-

The toxicity (Tox) is expressed as the pCC₅₀ value as determined with mock transfected cells while the pEC₅₀ for the wild type is displayed in column WT.

5 **Table 3.** Results of the toxicity testing and the resistance testing against strain A to F (expressed as pEC₅₀).

Compound	Strain A	Strain B	Strain C	Strain D	Strain E	Strain F	Tox	WT
1	8.53	8.44	8.38	8.65	8.51	ND	4.16	8.26
2	8.68	8.59	8.54	8.69	8.50	8.45	4.07	8.18
3	7.52	8.05	7.81	7.44	7.66	7.27	4.13	8.34
4	8.44	8.93	8.93	8.93	8.89	8.06	<5	9.34
5	9.37	9.57	9.71	ND	ND	8.71	4.15	9.26
6	6.70	6.89	7.47	6.96	ND	6.15	<4	8.4
7	6.37	7.53	7.49	6.93	7.36	6.11	4.33	8.23
8	7.5	7.87	7.59	7.47	7.56	6.85	<5	8.18
9	6.58	8.25	5.31	7.38	7.62	ND	4.29	8.31
10	7.07	8.03	7.80	7.64	7.88	7.06	4.14	8.04
11	6.95	8.14	8.12	8.08	8.14	6.99	4.24	7.84
12	6.64	8.12	6.72	7.58	8.11	ND	<4	8.37
13	7.39	8.24	8.42	8.13	8.57	6.98	<4	8.52
14	6.05	7.57	6.75	7.40	7.52	ND	4.33	8.42
15	7.29	7.54	7.40	7.30	7.44	6.64	4.04	7.95
16	<4	<4	<4	<4	<4	<4	4.95	5.85
17	7.50	8.18	7.91	7.63	8.12	6.80	4.2	8.15
18	7.51	8.21	8.13	7.67	8.04	6.83	4.07	8.72
19	7.36	7.80	7.88	7.43	7.90	6.80	ND	8.51
20	6.50	7.61	7.40	7.38	7.59	6.12	<4	8.32
22	6.93	7.83	8.19	7.78	8.36	6.1	<4	8.84
23	6.54	8.02	8.06	7.67	8.14	5.20	4.16	8.34
25	7.36	7.70	7.75	7.39	7.76	6.32	4.85	8.39
26	7.52	8.40	8.14	8.08	8.21	7.25	ND	8.57
27	6.80	7.69	5.30	7.07	7.51	6.19	<4	7.72
28	7.71	8.25	8.21	7.43	8.15	7.11	ND	8.60

ND means not determined

10 Some compounds have been tested for an even broader range of mutant HIV protease viruses. For instance, compound 1 was tested against a panel of more than 20 mutant

proteases whereby compound 1 had a pIC_{50} value of 9.13 for the most sensitive mutant and a pIC_{50} value of 8.12 for the most resistant mutant. This indicates that all the mutants within this set of more than 20 mutant proteases are sensitive within a narrow window of IC_{50} values and thus also in fold resistance values.

5

Table 4 : fold resistance

Compound	Strain A	Strain B	Strain C	Strain D	Strain E	Strain F
1	0.5	0.7	0.8	0.4	0.6	-
2	0.3	0.4	0.4	0.3	0.5	0.5
3	6.6	1.9	3.4	7.9	4.8	11.7
4	7.9	2.6	2.6	2.6	2.8	19.1
5	0.8	0.5	0.4	-	-	3.5
6	50.1	32.4	8.5	27.5	-	177.8
7	72.4	5.0	5.5	20.0	7.4	131.8
8	4.8	2.0	3.9	5.1	4.2	21.4
9	53.7	1.1	1000.0	8.5	4.9	-
10	9.3	1.0	1.7	2.5	1.4	9.5
11	7.8	0.5	0.5	0.6	0.5	7.1
12	53.7	1.8	44.7	6.2	1.8	-
13	13.5	1.9	1.3	2.5	0.9	34.7
14	234.4	7.1	46.8	10.5	7.9	-
15	4.6	2.6	3.5	4.5	3.2	20.4
17	4.5	0.9	1.7	3.3	1.1	22.4
18	16.2	3.2	3.9	11.2	4.8	77.6
19	14.1	5.1	4.3	12.0	4.1	51.3
20	66.1	5.1	8.3	8.7	5.4	158.5
22	81.3	10.2	4.5	11.5	3.0	549.5
23	63.1	2.1	1.9	4.7	1.6	1380.4
25	10.7	4.9	4.4	10.0	4.3	117.5
26	11.2	1.5	2.7	3.1	2.3	20.9
27	8.3	1.1	263.0	4.5	1.6	33.9
28	7.8	2.2	2.5	14.8	2.8	30.9

Protein Binding analyses:

Human serum proteins like albumin (HSA) or alpha-1 acid glycoprotein (AAG) are known to bind many drugs, resulting in a possible decrease in the effectiveness of the

drug. In order to determine whether the present compounds would be adversely effected by this binding, the anti-HIV activity of some of the present compounds was measured in the presence of human serum, thus evaluating the effect of the binding of the protease inhibitors to those proteins.

- 5 MT4 cells are infected with HIV-1 LAI at a multiplicity of infection (MOI) of 0.001-0.01 CCID₅₀ (50% cell culture infective dose per cell, CCID₅₀). After 1 h incubation, cells are washed and plated into a 96 well plate containing serial dilutions of the compound in the presence of 10% FCS (foetal calf serum), 10% FCS + 1 mg/ml AAG (α_1 -acid glycoprotein), 10% FCS + 45 mg/ml HSA (human serum albumin) or 50%
 10 human serum (HS). After 5 or 6 days incubation, the EC₅₀ (50% effective concentration in cell-based assays) is calculated by determining the cell viability or by quantifying the level of HIV replication. Cell viability is measured using the assay described above. Into a 96 well plate containing serial dilutions of the compound in the presence of 10% FCS or 10% FCS + 1 mg/ml AAG, HIV (wild type or resistant strain) and MT4 cells are
 15 added to a final concentration of 200-250 CCID₅₀/well and 30,000 cells/well, respectively. After 5 days of incubation (37°C, 5% CO₂), the viability of the cells is determined by the tetrazolium colorimetric MTT (3-[4,5-Dimethylthiazol-2-yl]-2,5-di-phenyltetrazolium bromide) method (Pauwels et al. J Virol. Methods 1988, 20, 309321).

20 Pharmacokinetic data

- The pharmacokinetic properties of some compounds of formula (I) were tested on rats and dogs. The compounds were evaluated in Whistar rats, source Iffa Credo, weighing approximately 350 g. Before dosing the animals were fasted overnight (approximately 12 h fasting period). The compounds were dissolved in DMSO or PEG 400. The
 25 results represented in the table concern the results from the oral or intra-peritoneal dosing of the compounds. Blood was sampled at 30 min, 1h, 2h, 3h, no pre-dose sample was taken. The amount of the compound in the biological sample was determined using LC-MS. In the table below "or" means oral dosing, "ip" means intra-peritoneal dosing, "mpk" means mg per kilogram. The results are illustrated in Table 5.

30

Table 5

Compound	C _{max} (ng/ml)	conditions (results normalized to 10mpk)
2	427 after 30 minutes	ip, rat, DMSO
2	52 after 30 minutes	or, rat, PEG400
4	1668 after 30 minutes	ip, rat, DMSO
4	348 after 30 minutes	or, rat, DMSO
4	225 after 30 minutes	or, rat, PEG400
15	86 after 30 minutes	ip, rat, DMSO
15	10 after 180 minutes	or, rat, PEG400

18	1141 after 240 minutes	ip, rat, DMSO
18	396 after 30 minutes	or, rat, DMSO
18	668 after 15 minutes	or, rat, PEG400
18	15 after 60 minutes	or, dog, DMSO
18	42 after 30 minutes	or, dog, PEG400
21	1763 after 15 minutes	ip, rat, DMSO
21	1139 after 15 minutes	or, rat, DMSO
21	1315 after 15 minutes	or, rat, PEG400
21	61 after 120 minutes	or, dog, PEG400 – 2 animals
21	184 after 30 minutes	or, dog, PEG400 – 4 animals
25	453 after 30 minutes	ip, rat, DMSO
28	1003 after 30 minutes	ip, rat, DMSO
28	540 after 30 minutes	or, rat, DMSO
28	430 after 60 minutes	or, rat, PEG400

Formulation

Active ingredient, *in casu* a compound of formula (I), is dissolved in organic solvent such as ethanol, methanol or methylene chloride, preferably, a mixture of ethanol and methylene chloride. Polymers such as polyvinylpyrrolidone copolymer with vinyl acetate (PVP-VA) or hydroxypropylmethylcellulose (HPMC), typically 5 mPa.s, are dissolved in organic solvents such as ethanol, methanol methylene chloride. Suitably the polymer is dissolved in ethanol. The polymer and compound solutions are mixed and subsequently spray dried. The ratio of compound/polymer can be selected from 1/1 to 1/6. Intermediate ranges were 1/1.5 and 1/3. The spraydried powder, a solid dispersion, is subsequently filled in capsules for administration. The drug load in one capsule depends on the capsule size used.

Film-coated Tablets

Preparation of Tablet Core

A mixture of 100 g of active ingredient, *in casu* a compound of formula (I), 570 g lactose and 200 g starch was mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture was sieved, dried and sieved again. Then there was added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole was mixed well and compressed into tablets, giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating


To a solution of 10 g methylcellulose in 75 ml of denaturated ethanol there was added a solution of 5 g of ethylcellulose in 150 ml of dichloromethane. Then there were added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene glycol was molten and dissolved in 75 ml of dichloromethane. The latter solution was added

-35-

to the former and then there were added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated color suspension and the whole was homogenated. The tablet cores were coated with the thus obtained mixture in a coating apparatus.

CLAIMS

1. The use of a 2-amino-benzothiazole, having the formula



a *N*-oxide, salt, stereoisomeric form, racemic mixture, prodrug, ester or metabolite thereof, wherein

R_1 is hexahydrofuro[2,3-*b*]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C_{1-6} alkyl, hydroxy, amino, halogen, amino C_{1-4} alkyl and mono- or di(C_{1-4} alkyl)amino;

10 R_2 is hydrogen or C_{1-6} alkyl;

L is a direct bond, -O-, C_{1-6} alkanediyl-O- or -O- C_{1-6} alkanediyl;

R_3 is phenyl C_{1-4} alkyl;

R_4 is C_{1-6} alkyl;

R_5 is hydrogen or C_{1-6} alkyl;

15 R_6 is hydrogen or C_{1-6} alkyl;

in the manufacture of a medicament useful for inhibiting mutant HIV protease in a mammal infected with said mutant HIV protease.

2. The use according to claim 1 wherein

20 R^2 is hydrogen;

R^3 is phenylmethyl;

R^4 is C_{1-4} alkyl, preferably isobutyl;

R^5 is hydrogen or methyl;

R^6 is hydrogen or methyl.

25

3. The use according to claim 1 or 2 wherein R^5 is methyl or hydrogen and R^6 is hydrogen

4. The use according to any one of claims 1 to 3 wherein both R^5 and R^6 are hydrogen.

30

5. The use according to any one of claims 1 to 4 wherein - L - R^1 is -O- (hexahydrofuro[2,3-*b*]furanyl), -O-tetrahydrofuranyl, -O-methyl-(optionally substituted phenyl), -O-methyl-pyridinyl, -O-methyl-thiazolyl, -O-methyl-thiazolyl, -methyl-O- (optionally substituted phenyl) or optionally substituted phenyl.

35

BEST AVAILABLE COPY

6. The use according to any one of claims 1 to 5 wherein the compound is
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
5 carbamic acid thiazol-5-ylmethyl ester;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxy-
propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
10 {3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid benzyl ester;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
15 carbamic acid pyridin-3-ylmethyl ester;
3-amino-N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-3-hydroxy-2-methyl-benzamide;
20 {3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid tetrahydro-furan-3-yl ester;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-methyl-benzamide;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
25 propyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-3-fluoro-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-(4-aminomethyl-2,6-dimethyl-phenoxy)-acetamide;
30 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-carbamic acid thiazol-5-ylmethyl ester;
3-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-
sulfonyl)-amino]-propyl}-2-methyl-benzamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
35 propyl}-carbamic acid tetrahydro-furan-3-yl ester;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-3-hydroxy-2-methyl-benzamide;


BEST AVAILABLE COPY

- N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-iodo-2,6-dimethyl-phenoxy)-acetamide;
 2-(4-aminomethyl-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
 5 2-(4-amino-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
 N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-4-bromo-2-methyl-benzamide;
 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
 10 propyl}-carbamic acid oxazol-5-ylmethyl ester;
 4-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide;
 or a salt, or a stereoisomeric form thereof.

- 15 7. The use according to any one of claims 1 to 6 wherein the mutant HIV protease has at least one mutation at a position selected from 10, 71 and 84.

8. The use according to any one of claims 1 to 7 wherein the fold resistance of the mutant HIV protease for the compound described in any one of claims 1 to 6 ranges
 20 between 0.01 and 100.

9. A compound having the formula



- a *N*-oxide, salt, stereoisomeric form, racemic mixture, prodrug, ester or metabolite
 25 thereof, wherein
 R₁ is hexahydrofuro[2,3-*b*]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C₁₋₆alkyl, hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono- or di(C₁₋₄alkyl)amino;
 30 R₂ is hydrogen or C₁₋₆alkyl;
 L is a direct bond, -O-, C₁₋₆alkanediyl-O- or -O-C₁₋₆alkanediyl;
 R₃ is phenylC₁₋₄alkyl;
 R₄ is C₁₋₆alkyl;
 R₅ is hydrogen or C₁₋₆alkyl;
 35 R₆ is hydrogen or C₁₋₆alkyl;

BEST AVAILABLE COPY

provided that the compound is other than :

- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid benzyl ester;
- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid pyridin-3-ylmethyl ester;
- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid thiazol-5-ylmethyl ester;
- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
- 3-amino-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
- 4-amino-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
- 5-amino-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
- N-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
- N-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-4-hydroxy-2-methyl-benzamide;
- N-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-hydroxy-2-methyl-benzamide; and
- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (S)-(tetrahydrofuran-3-yl) ester.

10. A compound according to claim 9 wherein R¹ is hexahydrofuro[2,3-b]furanyl or oxazolyl.

11 A compound according to claim 9 wherein R₁ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, and L is a direct bond.

30

12. A compound according to claim 9 wherein R₁ is hexahydrofuro[2,3-b]furanyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C₁₋₆alkyl, hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono-or di(C₁₋₄alkyl)amino; and L is -O-.

35

13. A compound according to claim 9 wherein R₁ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, or phenyl substituted with one or more substituents independently selected from C₁₋₆alkyl, hydroxy, amino, halogen, aminoC₁₋₄alkyl and

mono-or di(C₁₋₄alkyl)amino; and L is C₁₋₆alkanediyl-O- whereby the -O- is attached to the nitrogen of the amide.

14. A compound according to claim 9 wherein R₁ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono-or di(C₁₋₄alkyl)amino; and L is -O-C₁₋₆alkanediyl whereby -O- is attached to the R¹ group.
15. A compound according to any one of claims 9 to 14 wherein at least one of R₅ and R₆ is C₁₋₆alkyl.
16. A compound according to any one of claims 9 to 15 wherein R² is hydrogen; R³ is phenylmethyl; R⁴ is C₁₋₄alkyl.
17. a compound according to any one of claims 9 to 16 having the formula
 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
 {1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxy-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
 N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-fluoro-2-methyl-benzamide;
 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-aminomethyl-2,6-dimethyl-phenoxy)-acetamide;
 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid thiazol-5-ylmethyl ester;
 3-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-2-methyl-benzamide;
 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid tetrahydro-furan-3-yl ester;
 N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide;
 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(4-iodo-2,6-dimethyl-phenoxy)-acetamide;
 2-(4-aminomethyl-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;

- 2-(4-amino-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-4-bromo-2-methyl-benzamide;
5 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid oxazol-5-ylmethyl ester;
4-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide;
a salt thereof, or a stereoisomeric form thereof.
- 10 18. A compound according to claim 9 or 10 having the formula
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
15 {1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
or a salt or stereoisomeric form thereof.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/425 C07D277/82 C07D493/04 C07D417/12 A61P31/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 06030 A (MUELLER RICHARD A ;VAZQUEZ MICHAEL L (US); MONSANTO CO (US); SEARL) 2 March 1995 (1995-03-02) cited in the application page 195	1-18
A	WO 99 67254 A (UNIV ILLINOIS ;US HEALTH (US); ERICKSON JOHN W (US); GHOSH ARUN K) 29 December 1999 (1999-12-29) cited in the application the whole document	1-18
P,X	WO 02 083657 A (DE BETHUNE MARIE-PIERRE ;DE KERPEL JAN OCTAAF ANTOON (BE); DE KOCK) 24 October 2002 (2002-10-24) * see example 4, intermediate a-8; p 26, lines 20-22 *	1-18

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

13 October 2003

Date of mailing of the international search report

29/10/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bérillon, L

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9506030	A	02-03-1995	US 5843946 A	01-12-1998
			US 6060476 A	09-05-2000
			AT 174587 T	15-01-1999
			AU 7669794 A	21-03-1995
			DE 69415326 D1	28-01-1999
			DE 69415326 T2	02-06-1999
			DK 715618 T3	23-08-1999
			EP 0715618 A1	12-06-1996
			ES 2127938 T3	01-05-1999
			GR 3029378 T3	28-05-1999
			WO 9506030 A1	02-03-1995
			US 6248775 B1	19-06-2001
			US 5830897 A	03-11-1998
			US 6172082 B1	09-01-2001
			US 5786483 A	28-07-1998
			US 6335460 B1	01-01-2002
			US 6472407 B1	29-10-2002
			US 6534493 B1	18-03-2003
			US 2002052399 A1	02-05-2002
			US 5744481 A	28-04-1998
			US 5968942 A	19-10-1999
			US 6500832 B1	31-12-2002
WO 9967254	A	29-12-1999	AU 4828099 A	10-01-2000
			AU 4828199 A	10-01-2000
			CA 2336160 A1	29-12-1999
			EP 1088098 A2	04-04-2001
			JP 2002518063 T	25-06-2002
			WO 9967417 A2	29-12-1999
			WO 9967254 A2	29-12-1999
WO 02083657	A	24-10-2002	WO 02083657 A2	24-10-2002

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.